

## Egg yolk (emulsion)

Used as a supplement in many culture media for the isolation or identification of *Bacillus* and *Staphylococcus*.

► Storage : 8 - 14 °C

Cat. No.	Description	Pack.	€ Excl. VAT
77780	100 ml bottle of egg yolk emulsion - Condalab	1	NC -

## Egg yolk with potassium tellurite

For the preparation of the Baird Parker according to ISO 6888.

► Storage : 8 - 14 °C

Cat. No.	Description	Pack.	€ Excl. VAT
777800	100 ml bottle of egg yolk emulsion with potassium telluric - Condalab	1	NC -

## KAA presumption (broth)

For the presumption and detection of intestinal enterococci in foods according to Mossel.

► Storage : 2 - 25 °C

► Incubation : 35 ± 2 °C for 24 - 48 h

Cat. No.	Description	Pack.	€ Excl. VAT
777267	Dehydrated medium KAA presumption broth - Condalab	500 g	NC -

## KAA confirmation (agar)



For the isolation and confirmation of intestinal enterococci in foods according to MOSEL.

► Storage : 2 - 25 °C

► Incubation : 35 ± 2 °C for 24 - 48 h

*Enterococcus faecalis*

Cat. No.	Description	Pack.	€ Excl. VAT
777266	Dehydrated medium KAA confirmation agar - Condalab	500 g	NC -

## Kligler Hajna (agar)

For the presumptive identification of enterobacteria based on the fermentation of glucose, lactose, sucrose and the production of gas and H<sub>2</sub>S.

For the detection of salmonella in food and dairy products.

► Storage : 2 - 25 °C in the dark

► Incubation : 37 °C for 18 - 24 h

Cat. No.	Description	Pack.	€ Excl. VAT
693436	Kligler Hajna Agar Slant Tube - ClearLine®	100	NC -

## Kligler iron (agar)

Medium for the detection of yersinia enterocolitica.

► Storage : 2 - 25 °C

► Incubation : 30 °C for 18 - 24 h

Cat. No.	Description	Pack.	€ Excl. VAT
777275	Dehydrated medium Kligler iron agar - Condalab	500 g	NC -

## Koser citrate (broth)

For the differentiation of *Escherichia coli* from enterobacteria based on the use of citrate.

► Storage : 2 - 25 °C

► Incubation : 35 ± 2 °C for 18 - 24 h

Cat. No.	Description	Pack.	€ Excl. VAT
777276	Dehydrated medium Koser citrate broth - Condalab	500 g	NC -

## Lactose (broth)

For the culture of coliforms and salmonella in water, food, dairy products.

► Storage : 2 - 25 °C

► Incubation : 35 ± 2 °C for 18 - 24 h

Cat. No.	Description	Pack.	€ Excl. VAT
777278	Dehydrated medium lactose broth - Condalab	500 g	NC -

## Lactose sulphite (broth)

For the culture of coliforms and salmonella in water, food, dairy products.

► Storage : 2 - 25 °C for the dehydrated base and 2 - 8 °C for the ready-to-use tubes

► Incubation : 46 °C for 18 - 24 h

### Dehydrated medium

Cat. No.	Description	Pack.	€ Excl. VAT
777279	Dehydrated medium for lactose sulphite broth - Condalab	500 g	NC -

### Supplement

► After heating the tube to 100 °C and allowing it to cool for 5 minutes, add 0,5 ml of a 12 g/l solution of sodium metabisulphite and 0,5 ml of a 10 g/l solution of ammoniacal ferric citrate sterilised by filtration through a 0,45 µm membrane

### Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693314	8 ml tube of Durham broth lactose sulphite - ClearLine®	100	NC -
693408	9 ml tube with Durham bell broth lactose sulphite - ClearLine®	100	NC -

## Lauryl sulphate (broth) - Lauryl Tryptose Broth (LTB)

Used to detect coliforms in food, water, milk and dairy products.

The use of inoculum larger than 1 ml requires the use of double strength lauryl sulphate broth to ensure that sufficient nutrients are maintained.

► Storage : 2 - 25 °C

► According to ISO 4831 : 30 °C ou 37 °C for 24 h

► According to ISO 7251 : 37 °C for 48 h, examining gas production after 24 h and 48 h

Cat. No.	Description	Pack.	€ Excl. VAT
777281	Dehydrated medium for lauryl sulfate broth - Condalab	500 g	NC -

## Lauryl sulphate (agar)

Selective medium for the detection of coliforms in water, dairy products and seafood.

► Storage : 2 - 25 °C

► Incubation : 35 ± 2 °C for 24 - 48 h

Cat. No.	Description	Pack.	€ Excl. VAT
777280	Dehydrated medium for lauryl sulfate agar - Condalab	500 g	NC -

## Chromogenic Lauryl Sulphate (broth)

Enrichment medium for the simultaneous detection of total coliforms and *Escherichia coli* in water, food, dairy products.

► Storage : 2 °C - 8 °C

► Incubation : 35 °C ± 2 °C for 18 - 24 h. Control of the tubes under UV 366 nm lamp. Blue light indicates the presence of *E. coli*

Cat. No.	Description	Pack.	€ Excl. VAT
777848	Dehydrated medium for lauryl sulphate chromogenic broth - Condalab	500 g	NC -