

Acetate differential agar

See Acetate differential agar

Actinomycete isolation agar glycerol

See actinomycete isolation agar

American bacteriological agar

Gelling agent for culture media.

- Storage : 2 - 25 °C
- Gel strength : 600 - 850 g/cm²

Cat. No.	Description	Pack.	€ Excl. VAT
777475	American bacteriological agar - Condalab	500 g	NC -

European bacteriological agar

Gelling agent for culture media.

- Storage : 2 - 25 °C
- Gel strength : 600 - 1100 g/cm²

Cat. No.	Description	Pack.	€ Excl. VAT
777476	European bacteriological agar Condalab	500 g	NC -
777525	Condalab European bacteriological agar	5 kg	NC -

Industrial agar

Gelling agent for culture media.

- Storage : 2 - 25 °C
- Gel strength : 850 g/cm²

Cat. No.	Description	Pack.	€ Excl. VAT
777477	Condalab Industrial Agar	500 g	NC -

Agar LB (Lennox)

See LB agar

Agar Luria

See Luria agar

Pharmaceutical agar

Gelling agent for culture media.

- Storage : 2 - 25 °C
- Gel strength : 750 - 1000 g/cm²

Cat. No.	Description	Pack.	€ Excl. VAT
777479	Condalab Pharmaceutical Agar	500 g	NC -

Purified agar

Gelling agent for culture media.

- Storage : 2 - 25 °C
- Gel strength : 700 - 1200 g/cm²

Cat. No.	Description	Pack.	€ Excl. VAT
777480	Condalab Purified agar	500 g	NC -

Agar for plant propagation

Gelling agent for culture media.

- Storage : 2 - 25 °C
- Gel strength : 900 g/cm²

Cat. No.	Description	Pack.	€ Excl. VAT
777478	Agar for plant propagation - Condalab	500 g	NC -

Agar YPD

See YPD agar

Anaerobic agar

See Anaerobic agar

Bacillus cereus (selective agar)

See mossel (Bacillus cereus agar)

Baird-Parker

Recommended for the detection and enumeration of coagulase-positive staphylococci, including *Staphylococcus aureus* in food.

- Storage : 2 - 25 °C (for dehydrated media) and 2 - 8 °C (for ready-to-use media)
- Incubation : 35 ± 2 °C for 24 - 48 h
- Incubation for ISO 11133 : 37 ± 1 °C for 24 ± 2 h then 24 ± 2 h more

Dehydrated bases

Cat. No.	Description	Pack.	€ Excl. VAT
777184	Baird-Parker agar dehydrated medium - Condalab	500 g	NC -

Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693396	ClearLine® Baird-Parker Agar - Flask 90 ml	10	NC -
693350	ClearLine® Baird-Parker Agar - Flask 100 ml	10	NC -

Supplement: egg yolk with tellurite (ISO 6888-1)

- Storage 8 - 14 °C

➤ Composition :

Potassium tellurite	2.1 g	Sodium chloride	4.25 g
Egg yolk	200 ml	Distilled water	800 ml

Cat. No.	Description	Pack.	€ Excl. VAT
777800	Egg yolk 100 ml bottle - Condalab	1	NC -

Supplement: RPF (ISO 6888-2)

- Storage : 8 - 14 °C

➤ Composition :

Potassium tellurite	2.5 mg	Trypsin	2.5 mg
Rabbit plasma	2.5 ml	Bovine fibrogen	0.38 g

Cat. No.	Description	Pack.	€ Excl. VAT
777523	RPF Supplement - Condalab	10 bottles	NC -

Nitrogen base for amino acid-free yeast

For the classification of yeasts according to their amino acid and carbohydrate requirements.

- Storage : 2 - 25 °C
- Incubation: 25 - 30°C for 2 - 5 days

Cat. No.	Description	Pack.	€ Excl. VAT
777508	Dehydrated nitrogen-based medium for amino acid-free yeast - Condalab	500 g	NC -