

Rogosa SL (Broth)

For the selection and cultivation of lactobacilli.

- **Storage :** 2 - 25 °C
- **Incubation :** 35 ± 2 °C for 18 - 48 h

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|--|-------|-------------|
| 777361 | Dehydrated Medium Rogosa SL broth - Condalab | 500 g | NC - |

Rogosa SL (Agar)

For the selection and cultivation of lactobacilli.

- **Storage :** 2 - 25 °C
- **Incubation :** 35 °C ± 2 °C for 24 - 48 h

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|---|-------|-------------|
| 777360 | Dehydrated Medium Rogosa SL agar - Condalab | 500 g | NC - |

Rose Bengal + Chloramphenicol (Agar)

A medium for the enumeration of yeasts and moulds in foods.

- **Storage :** 2 - 25 °C
- **Incubation :** 25 - 30 °C for 7 days

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|---|-------|-------------|
| 777362 | Dehydrated medium Rose Bengal agar + chloramphenicol - Condalab | 500 g | NC - |

Rose Bengal + Chloramphenicol + Dichloran ISO 11133 / ISO 21527 (DRBC - Agar)

A medium for the enumeration of yeasts and moulds in foods.

- **Storage :** 2 - 25 °C
- **Incubation :** 25 ± 1 °C for 5 days

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|---|-------|-------------|
| 777363 | Dehydrated medium Rose Bengal agar + chloramphenicol + Dichloran - Condalab | 500 g | NC - |
| 693360 | DRBC Agar 100 ml bottle - ClearLine® | 10 | NC - |



Aspergillus brasiliensis

Rothe (Broth)

Used for the enumeration of faecal enterococci.

- **Storage :** 2 - 25 °C
- **Incubation :** 35 ± 2 °C for 24 - 48 h

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|---|-------|-------------|
| 777364 | Dehydrated medium Rothe broth (Glucose broth with azide) - Condalab | 500 g | NC - |

Glucose-rich sabouraud (Broth)

For the growth of a wide variety of yeasts and moulds.

- **Storage :** 2 - 25 °C (dehydrated base and ready-to-use medium)
- **Incubation :** 20 - 25 °C for 2 to 5 days

Dehydrated base

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|---|-------|-------------|
| 777371 | Dehydrated medium sabouraud glucose agar - Condalab | 500 g | NC - |

Ready-to-use media

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|--|-------|-------------|
| 693416 | 10 ml tube of sabouraud glucosed broth - ClearLine® | 100 | NC - |
| 693381 | 100 ml bottle of sabouraud glucosed broth - ClearLine® | 10 | NC - |

Sabouraud glucose (Agar)

For the growth and isolation of a wide variety of yeasts and moulds.

- **Storage :** 2 - 25 °C (dehydrated base and ready-to-use medium)
- **Incubation :** 20 - 25 °C for 2 to 5 days

Dehydrated base

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|---|-------|-------------|
| 777366 | Dehydrated medium sabouraud glucose agar - Condalab | 500 g | NC - |

Ready-to-use media

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|--|-------|-------------|
| 693382 | 100 ml bottle of sabouraud glucose agar - ClearLine® | 10 | NC - |
| 693450 | 200 ml bottle of sabouraud glucose agar - ClearLine® | 6 | NC - |

Sabouraud + chloramphenicol (Agar)

For the growth and isolation of a wide variety of yeasts and moulds.

- **Storage :** 2 - 25 °C (dehydrated base and ready-to-use medium)
- **Incubation :** 20 - 25 °C for 2 to 5 days

Dehydrated base

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|--|-------|-------------|
| 777368 | Dehydrated medium glucosed sabouraud agar + chloramphenicol - Condalab | 500 g | NC - |

Ready-to-use media

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|--|-------|-------------|
| 693439 | Tube with Glucose sabouraud slant tube + chloramphenicol - ClearLine® | 100 | NC - |
| 693383 | 100 ml bottle of sabouraud glucose agar + chloramphenicol - ClearLine® | 10 | NC - |
| 693451 | 200 ml bottle of sabouraud glucose agar + chloramphenicol - ClearLine® | 6 | NC - |

Sabouraud + chloramphenicol + cycloheximide (Agar)

Allows the growth and isolation of a wide variety of yeasts and moulds. Chloramphenicol inhibits the growth of bacteria. Cycloheximide inhibits the growth of saprophytic fungi but has no effect on pathogenic fungi.

- **Storage :** 2 - 25 °C (dehydrated base) - 2 - 8 °C (ready-to-use medium)
- **Incubation :** 30 °C for 2-5 days

Dehydrated base

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|--|-------|-------------|
| 777370 | Dehydrated medium glucosed sabouraud agar + chloramphenicol + cycloheximide - Condalab | 500 g | NC - |

Ready-to-use media

| Cat. No. | Description | Pack. | € Excl. VAT |
|----------|---|-------|-------------|
| 693440 | Slant agar tube with sabouraud glucose + chloramphenicol + cycloheximide - ClearLine® | 100 | NC - |
| 693332 | 200 ml bottle sabouraud glucose agar + chloramphenicol + cycloheximide - ClearLine® | 6 | NC - |