

## T.S.B. (Trypticasein Soy Broth)

A general purpose medium, allowing the growth of a wide variety of micro-organisms. It is recommended by the European and American pharmacopoeia.

- **Storage :** 2 - 25 °C (Dehydrated base and ready-to-use medium)
- **Incubation :** 30 ± 1 °C for 48 ± 4 h

### Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777413	Trypticasein Soy Broth (TSB) - Condalab	500 g	NC -

### Ready-to-use media

Cat. No.	Description	Units/box	€ Excl. VAT/box
693419	10 ml tube with Tryptic soy broth (TSB) - ClearLine®	100	NC -
693404	100 ml bottle with Tryptic soy broth (TSB) - ClearLine®	10	NC -
693454	200 ml bottle with Tryptic soy broth (TSB) - ClearLine®	6	NC -

## T.S.C. (Tryptose sulphite cycloserine - Agar)

Recommended for the detection and enumeration of Clostridium perfringens and sulphite-reducing anaerobes in water and food products.

- **Storage :** 2 - 25°C (dehydrated base) - 2 - 8°C (ready-to-use medium)
- **Incubation :** 37 °C for 20 h

### Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777424	Dehydrated medium T.S.C. - Condalab	500 g	NC -

### Supplement

For 500 ml of medium.

Cat. No.	Description	Pack.	€ Excl. VAT
777524	Supplement D - Cycloserine for T.S.C. agar - Condalab	10 bottles for 500 ml	NC -

## T.S.I. (Triple Sugar Iron - Agar) ISO 19250



Escherichia coli

Used for presumptive identification of Enterobacteriaceae based on fermentation of glucose, lactose, sucrose and the production of gas and H<sub>2</sub>S. Its use is recommended for the detection of salmonella in pharmaceuticals, and for the detection of salmonella and campylobacter in food.

- **Storage :** 2 - 25 °C
- **Incubation :** 36 ± 2 °C for 24 ± 3 h

Cat. No.	Description	Pack.	€ Excl. VAT
777407	Dehydrated medium T.S.I. agar - Condalab	500 g	NC -

## T.S.N. (Tryptone sulfite with neomycin)

Recommended for the detection and enumeration of Clostridium perfringens and sulphite reducing anaerobes in certain foods.

- **Storage :** 2 - 25°C (dehydrated base) and 2 - 8°C (ready-to-use medium)
- **Incubation :** 48 ± 1 °C for 18 - 24 h

### Dehydrated base

Cat. No.	Description	Pack.	€ Excl. VAT
777416	Dehydrated medium T.S.N. agar - Condalab	500 g	NC -

### Ready-to-use media

Cat. No.	Description	Pack.	€ Excl. VAT
693420	20 ml tube of T.S.N. agar - ClearLine®	100	NC -

## Tryptone

Used as a source of nitrogen.

- **Storage :** 2 - 25 °C

Cat. No.	Description	Pack.	€ Excl. VAT
777472	Tryptone - Condalab	500 g	NC -

## Tryptone salt

See Peptone salt page 581

## Tryptose sulphite

For the detection of clostridium perfringens.

- **Storage :** 2 - 25 °C
- **Incubation :** 37 ± 1 °C for 20 ± 4 h; 44 ± 4 h

Cat. No.	Description	Pack.	€ Excl. VAT
777422	Dehydrated Medium Tryptone Sulphite Agar - Condalab	500 g	NC -

## T.S.Y.E.A. Agar (Tryptone Soy Yeast Extract Agar)

For confirmation of Listeria spp

- **Storage :** 2 - 25 °C
- **Incubation :** 37 °C for 18 - 24 h

Cat. No.	Description	Pack.	€ Excl. VAT
777416	Dehydrated medium for T.S.Y.E.A agar - Condalab	500 g	NC -

## T.S.Y.E.B. (Tryptone Soy Broth Yeast Extract)

For confirmation of Listeria monocytogenes.

- **Storage :** 2 - 25 °C
- **Incubation :** 25 °C for 18 - 24 h

Cat. No.	Description	Pack.	€ Excl. VAT
777417	Dehydrated medium for T.S.Y.E.B broth - Condalab	500 g	NC -

## U.B.A. (Universal beer Agar) (Agar)

For the cultivation of important bacteria in the beer industry.

- **Storage :** 2 - 25 °C
- **Incubation :** 28 - 30 °C for 3 days

Cat. No.	Description	Pack.	€ Excl. VAT
777888	Dehydrated medium for U.B.A. agar - Condalab	500 g	NC -